



available at www.sciencedirect.com



journal homepage: www.elsevierhealth.com/journals/ctim



A detailed safety assessment of a saw palmetto extract^{☆,☆☆}

Andrew L. Avins^{a,b,c,d,*}, Stephen Bent^{b,c,e}, Suzanne Staccone^b, Evelyn Badua^f, Amy Padula^b, Harley Goldberg^a, John Neuhaus^d, Esther Hudes^d, Katusto Shinohara^{f,g}, Christopher Kane^{f,g}

^a Division of Research, Northern California Kaiser Permanente, United States

^b General Internal Medicine Section, San Francisco VA Medical Center, United States

^c Department of Medicine, University of California, San Francisco, United States

^d Department of Epidemiology and Biostatistics, University of California, San Francisco, United States

^e Osher Center for Integrative Medicine, Department of Medicine, University of California, San Francisco, United States

^f Urology Section, San Francisco VA Medical Center, United States

^g Department of Urology, University of California, San Francisco, United States

Available online 20 February 2008

KEYWORDS

Saw palmetto;
Drug toxicity;
Benign prostatic
hyperplasia

Summary

Background: Saw palmetto is commonly used by men for lower-urinary tract symptoms. Despite its widespread use, very little is known about the potential toxicity of this dietary supplement.

Methods: The Saw palmetto for Treatment of Enlarged Prostates (STEP) study was a randomized clinical trial performed among 225 men with moderate-to-severe symptoms of benign prostatic hyperplasia, comparing a standardized extract of the saw palmetto berry (160 mg twice daily) with a placebo over a 1-year period. As part of this study, detailed data were collected on serious and non-serious adverse events, sexual functioning, and laboratory tests of blood and urine. Between-group differences were assessed with mixed-effects regression models.

Results: There were no significant differences observed between the saw palmetto and placebo-allocated participants in the risk of suffering at least one serious adverse event (5.4% vs. 9.7%, respectively; $p=0.31$) or non-serious symptomatic adverse event (34.8% vs. 30.1%, $p=0.48$). There were few significant between-group differences in sexual functioning or for most laboratory analyses, with only small differences observed in changes over time in total bilirubin ($p=0.001$), potassium ($p=0.03$), and the incidence of glycosuria (0% in the saw palmetto group vs. 3.7% in the placebo group, $p=0.05$).

[☆] This work was supported by a grant from the National Institute of Diabetes, Digestive, and Kidney Diseases (#R01 DK56199) and the National Center for Complementary and Alternative Medicine (#K08 AT001338).

^{☆☆} The study sponsors played no role in the study design; the collection, analysis, and interpretation of the data; in the writing of the manuscript; in the decision to submit a manuscript for publication.

* Corresponding author at: Division of Research, Northern California Kaiser Permanente, 2000 Broadway, 3rd Floor, Oakland, CA 94612, United States. Tel.: +1 510 891 3557; fax: +1 510 891 3606.

E-mail address: andrew.avins@ucsf.edu (A.L. Avins).

0965-2299/\$ – see front matter © 2007 Elsevier Ltd. All rights reserved.

doi:10.1016/j.ctim.2007.10.005

Conclusions: Despite careful assessment, no evidence for serious toxicity of saw palmetto was observed in this clinical trial. Given the sample size and length of this study, however, these data do not rule out potential rare adverse effects associated with the use of saw palmetto.

© 2007 Elsevier Ltd. All rights reserved.

Introduction

Herbal therapies are one of the most widely used alternative modalities in the U.S. with sales exceeding \$18 billion in 2005.¹ Among the most commonly used phytotherapeutics is an extract of the berry of the saw palmetto plant, a dwarf palm tree native to the southeastern U.S.^{2,3} Saw palmetto extracts are generally used to relieve symptoms associated with benign prostatic hyperplasia (BPH), a non-malignant enlargement of the prostate gland affecting the majority of men over the age of 50.⁴ Because saw palmetto extracts are sold without prescription, it is difficult to determine the numbers of men who take the extract regularly, but it is estimated that the number of regular users is approximately 2.5 million adults in the U.S.⁵

The efficacy of saw palmetto (also known as *Serenoa repens*) is currently the subject of active investigation by several research groups. An updated systematic review of saw palmetto for BPH found that most, but not all, published studies showed some modest benefit in overall lower-urinary tract symptoms and nocturia, but that much of the available research suffered from serious methodologic problems.⁶ We recently reported the efficacy results of a year-long clinical trial of a saw palmetto extract in men with at least moderately severe BPH that addressed many of the shortcomings of earlier studies. This trial, the Saw palmetto Trial for Enlarged Prostates (STEP) study, found no evidence of efficacy of saw palmetto for either BPH symptoms or objective measures of urinary function.⁷

Despite the substantial quantity of clinical research performed on saw palmetto, there are few data regarding potential adverse effects associated with its use. Most trials were of short duration; few described any systematic attempt to assess adverse effects, and with rare exception,⁸ laboratory testing was not performed to test for asymptomatic toxicities of saw palmetto.⁹ This information is of great public-health consequence given the large numbers of men who self-medicate with saw palmetto for extended periods of time. Since it is well known that most individuals who take dietary supplements do not inform their physicians about their use of these products,^{10–14} most men who take saw palmetto will not be monitored for potential adverse effects. Understanding the risks of using saw palmetto, therefore, is of great importance for patients, clinicians, and regulatory authorities.

A major goal of the STEP study was a detailed assessment of potential toxicity of saw palmetto, including both symptomatic adverse effects, as well as asymptomatic laboratory abnormalities. A summary of the major adverse-event data from this trial has been published previously.⁷ This report provides comprehensive information on the adverse-event data from the STEP trial, including detailed information about laboratory measurements.

Methods

Study design and participants

The STEP study was a single-center double-blind placebo-controlled randomized clinical trial of an extract of the saw palmetto berry. Inclusion criteria included age at least 50 years, a mean score of at least 8 on the American Urological Association Symptom Inventory (AUASI) on two measurements prior to randomization, a peak urine flow between 4 and 15 ml/s, a post-void residual volume <250 ml, and a prostate-specific antigen (PSA) <4.0 ng/ml (or a PSA <10 ng/ml with a negative prostate biopsy for malignancy). Potential participants were excluded if they had a creatinine >2.0 mg/dl, prior prostate surgery, a history of prostate cancer, a neurologic condition affecting urination, severe concomitant illness, or were taking a medication with androgenic or antiandrogenic properties. The STEP study was funded by the National Institutes of Health (co-funded by the National Institute of Diabetes, Digestive, and Kidney Diseases and the National Center for Complementary and Alternative Medicine) and was conducted under an Investigational New Drug exemption from the U.S. Food and Drug Administration. All study procedures were approved by the institutional review boards at the University of California, San Francisco and the Kaiser Foundation Research Institute.

Participants underwent two eligibility screening visits, a 1-month, single-blind run-in period, and were seen for follow-up visits at 1, 3, 6, 9, and 12 months after randomization.

Intervention

Participants were randomized to a saw palmetto berry extract, 160 mg twice daily, for 1 year or a placebo capsule. The saw palmetto preparation was produced by Indena, USA (Seattle, WA) and contained 92.1% total fatty acids. The extract was packaged in gelatin capsules by Cardinal Health (formerly RP Scherer Inc., St. Petersburg, FL) and supplied to the trial by Rexall-Sundown Inc., (Boca Raton, FL).

The identical-appearing placebo capsules contained 200 mg of polyethylene-glycol 400, colored to match the saw palmetto extract.

Outcomes

At each post-randomization visit, all participants were asked if they had experienced "any significant medical illness since the last study visit." Those who responded affirmatively to this global question were then asked to complete a symptom checklist that included open-ended fields.¹⁵ Serious adverse events (SAEs) were recorded and verified

Table 1 Baseline characteristics of study sample by treatment group

Characteristic	Saw palmetto (N = 112)	Placebo (N = 113)
Age (N (%))		
50–59 years	45 (40)	42 (37)
60–69 years	46 (41)	48 (42)
70–79 years	21 (18)	23 (20)
Race or ethnic group (N (%))		
White	94 (84)	89 (79)
Black	4 (4)	8 (7)
Asian or Pacific Islander	7 (6)	8 (7)
Hispanic	6 (5)	5 (4)
Other	1 (1)	2 (2)
American Urological Association Symptom Index		
Mean (S.D.)	15.7 (5.7)	15.0 (5.3)
Prostate volume (ml)		
Mean (S.D.)	34.7 (13.9)	33.9 (15.2)
Maximal urinary flow rate (ml/s)		
Mean (S.D.)	11.4 (3.5)	11.6 (4.3)
Post-void residual volume (ml)		
Mean (S.D.)	80.0 (51.9)	84.5 (63.8)

Abbreviations: N, number of participants; S.D., standard deviation.

with medical records, where possible. Non-serious adverse events were recorded and categorized by organ system.

Twenty-two laboratory tests were obtained at baseline and at 1, 6, and 12 months after randomization (Table 5). Most baseline laboratory values were obtained at the randomization visit, except the serum prostate-specific antigen test and the prothrombin time (international normalized ratio) were obtained at the first screening visit (approximately 6 weeks prior to randomization), as these were part of the eligibility screening process.

The effect of saw palmetto on sexual functioning was measured with the O’Leary Brief Sexual Function Inventory¹⁶ at randomization, the 6-month visit and at the 1-year close-out visit. Scores for each domain were calculated as the sum of scores for each of the items in that domain; each item was scored on a 0–4 Likert-like scale. The domains assessed (and the number of items included and the range of scores for that domain) were: sexual drive (two items, range 0–8), erectile function (three items, range 0–12), ejaculation (two items, range 0–8), perceptions of problems (three items, range 0–12), and overall sexual satisfaction (one item, range 0–4).

Statistical analyses

The frequencies of symptomatic adverse events (both serious and non-serious) were tabulated and the proportions of participants in each treatment group who reported at least one event were compared with Fisher’s exact tests.

Comparisons between the active-treatment and placebo arms for the laboratory and sexual functioning outcomes

were made with mixed-effects regression models to account for the repeated measures,¹⁷ which included a random intercept and terms to describe change in outcomes over time within each group. The test of the group-by-time interaction term provided the primary test of significance between the two treatment arms. This test assesses the statistical significance of the differences between the two groups for each continuous variable over the duration of the study. The time variable was treated as fully categorical (the individual time points (three for the sexual functioning outcome and four for the laboratory values) were modeled with indicator variables) as several variables showed significant departures from linear or quadratic models. The tables show the model-derived predicted mean values and standard errors for the baseline and closeout timepoints. Also shown are the differences between these assessments with confidence intervals derived using the standard error of the group-by-time (close-out value) interaction term from the mixed model, with the test of significance derived from the mixed-effects models. All analyses were consistent with the principle of intention-

Table 2 Summary of serious adverse events during STEP trial

SAE	ID	Adverse event	Treatment group
1	A	Hernia repair	Placebo
2	B	Hypotension	Placebo
3	B	Hematoma	Placebo
4	B	Bradycardia	Placebo
5	B	Coronary artery stent re-occlusion	Placebo
6	B	Coronary artery stent re-occlusion	Placebo
7	B	Superficial femoral artery occlusion	Placebo
8	B	Congestive heart failure	Placebo
9	C	Colon cancer	Placebo
10	D	Elective hip replacement	Placebo
11	E	Localized prostate cancer	Placebo
12	F	Total knee arthroplasty	Placebo
13	G	Syncope, possible seizure	Placebo
14	H	Gastrointestinal bleeding	Placebo
15	I	Shortness of breath	Placebo
16	J	Resection of bladder carcinoma	Placebo
17	J	Rhabdomyolysis	Placebo
18	K	Hip revision	Placebo
19	L	Lumbar laminectomy	Saw palmetto
20	L	Gastrointestinal bleeding	Saw palmetto
21	M	Vertigo	Saw palmetto
22	N	Bleeding gastric ulcer	Saw palmetto
23	O	Shoulder surgery	Saw palmetto
24	P	Atrial fibrillation	Saw palmetto
25	P	Elective laminectomy	Saw palmetto
26	Q	Melanoma removal	Saw palmetto

SAE: serious adverse event; ID: study participant identification code.

Table 3 Summary of non-serious adverse events during the STEP trial

Adverse event	Saw palmetto (N = 112)	Placebo (N = 113)
Cardiac		
Dysrhythmia/palpitations	0	2
Dermatologic		
Rash	1	3
Shingles	0	2
Skin cancer removal	1	0
Keratosis trunk	0	1
Gastrointestinal		
Diarrhea	2	2
Heartburn	0	3
Abdominal pain	2	1
Nausea/vomiting	2	0
Hemorrhoids	1	0
Abdominal swelling	1	0
Liver cyst on ultrasound scan	0	1
Polyp removal	1	0
Blood in stool	0	1
Genitourinary		
Nocturia	0	2
Discomfort in kidney	1	0
Pain in prostate area	1	0
Testicular pain	0	1
Kidney stone	1	0
Prostatitis	1	0
Urinary infection	1	0
HEENT		
Headache	1	0
TMJ pain	1	0
Head and neck infection	0	2
Chemical conjunctivitis	1	0
Periodontal cyst	1	0
Musculoskeletal		
Back pain	4	4
Gout	2	2
Joint pain/swelling	3	2
Trauma (fracture/bruise)	4	2
Myalgias	0	1
Soft tissue pain (e.g., tendonitis)	3	0
Infected digit	0	2
Neurologic/psychiatric		
Depression	1	0
Pulmonary		
Upper respiratory tract infection	12	10
Cough	1	2
Collapsed lung	1	0
Sleep apnea	1	0
Walking pneumonia	1	0
Miscellaneous/other		
Fatigue	0	2

Table 3 (Continued)

Adverse event	Saw palmetto (N = 112)	Placebo (N = 113)
Inguinal hernia	2	0
Axillary abscess	1	0
Cyst removal	0	1
Fistula	0	1
Hypothyroidism	0	2
Tumor removal	0	1
Yeast infection	1	0
Total non-serious AEs	57	53

AEs: adverse events.

to-treat in that all data for all participants were used; no data were imputed.

Results

Overall, 225 men were randomized to either the saw palmetto extract or a placebo; demographic characteristics were similar between the two groups (Table 1). Adherence to both the study visit schedule and medication regimen was excellent: 96% of randomized participants completed the study and 91.6% of all study medication was consumed (as measured by capsule counts at each visit);⁷ there was no difference in adherence between the two trial arms.

Serious adverse events

As described previously, a total of 26 serious adverse events among 17 participants were reported during the study (Table 2).⁷ While the majority of these events (18 events) occurred in participants randomized to placebo, several of these events occurred in the same participant, so that the likelihood of suffering at least one SAE was not significantly different between the two treatment arms: 5.4% in the saw palmetto group and 9.7% in the placebo group ($p=0.31$).

Most of the events were cardiovascular incidents, elective musculoskeletal procedures, or serious gastrointestinal problems. Three incident cancer cases occurred, of which two were in the placebo group. None of the serious adverse events were assessed as probably related to the study medicine and no deaths occurred.

Non-serious adverse events

The risk of suffering at least one non-serious adverse event was similar between the two groups (34.8% in the saw palmetto group and 30.1% in the placebo group, $p=0.48$). Events were distributed widely over many organ systems, with musculoskeletal, respiratory, and gastrointestinal problems being most commonly reported (Table 3).

Sexual functioning

No statistically significant differences were observed between the saw palmetto and placebo groups in the mea-

Table 4 Baseline and 12-month closeout scores for the domains of the O'Leary Brief Sexual Function Inventory

Domain (range)	Saw palmetto		Placebo		Difference	95% CI	p-Value ^a
	Baseline	12 months	Baseline	12 months			
Sexual drive (0–8)	4.29 ± 0.17	4.28 ± 0.18	4.26 ± 0.17	4.22 ± 0.18	0.03	−0.35 to 0.41	0.99
Erectile function (0–12)	5.43 ± 0.16	5.71 ± 0.16	5.42 ± 0.16	5.61 ± 0.16	0.08	−0.31 to 0.49	0.49
Ejaculation (0–8)	1.60 ± 0.22	1.55 ± 0.22	1.80 ± 0.22	1.90 ± 0.22	−0.16	−0.62 to 0.30	0.25
Perception of problems (0–12)	8.84 ± 0.33	8.79 ± 0.33	8.78 ± 0.33	8.38 ± 0.33	0.35	−0.31 to 1.01	0.04
Overall satisfaction (0–4)	2.18 ± 0.10	2.12 ± 0.11	2.01 ± 0.10	2.08 ± 0.10	−0.13	−0.40 to 0.14	0.40

Mean values at baseline and the 12-month closeout visit and between-group differences (calculated as the change in the placebo group subtracted from the change in the saw palmetto group). Mean values are derived from the mixed-effects regression models. The *p*-values for the between-group differences are derived from the hypothesis test on the group-by-time interaction term from the linear mixed-effects model which tests for overall differences between groups at any time point; the confidence interval for the between-group differences are obtained from the difference between the estimated baseline and closeout values and does not consider interim time points. CI: confidence interval.

^a *p*-Value for between-group differences in change in variable.

sured domains of sexual functioning with the exception of the perception-of-sexual-problems domain which showed a small but significantly greater improvement in the placebo group (Table 4).

Laboratory test results

A large number of serum laboratory tests were performed at baseline, 1 month, 6 months, and 1 year after randomization (Table 5). Only the between-group differences in total bilirubin and potassium achieved conventional levels of statistical significance (Table 5). The magnitude of the differences for each of these variables was judged to be small from a clinical standpoint. In addition, there were no significant differences in most urine tests, including pH ($p=0.50$), specific gravity ($p=0.37$), and the proportions of saw-palmetto-assigned participants vs. placebo-assigned participants whose closeout urine samples had evidence of hematuria (2.9% vs. 6.5%, $p=0.22$), proteinuria (1.0% vs. 4.6%, $p=0.11$), ketonuria (2.9% vs. 2.8%, $p=0.95$), or bilirubinuria (0% vs. 1.0%, $p=0.33$); patients randomized to placebo were significantly more likely to develop glycosuria (0% vs. 3.7%, $p=0.05$). Of note, we found no significant effect of saw palmetto on serum prostate-specific antigen levels, consistent with other studies.^{8,18–20}

Discussion

Because many men choose to take saw palmetto extracts, the potential adverse effects of this dietary supplement must be ascertained so that these individuals can make informed decisions about their use of this product. The STEP study provided a unique opportunity to make detailed assessments about potential toxicities of saw palmetto, having obtained extensive data on both symptomatic side effects as well as asymptomatic laboratory abnormalities and included a placebo group which allowed for comparison with an untreated control condition.

Overall, we found no evidence that consumption of this saw palmetto extract, at a dose of 160 mg twice daily over a period of 1 year, was associated with any clinically important adverse effects. Relatively few participants suf-

fered serious adverse events, and these were more common in the placebo-allocated participants. Non-serious adverse events were nearly equally distributed between the saw palmetto and placebo groups, both in total number and in the proportion of participants who suffered at least one adverse event. Only one of the five domains on the O'Leary sexual-functioning instrument (the perception-of-problems domain) showed a significant difference between treatment groups; however, this difference was small (approximately 1/3 of a point difference on a 12-point scale). Finally, we found little evidence of toxicity of saw palmetto among the laboratory analyses performed: while there were a small number of significant results, the large number of tests conducted would be expected to generate a small number of significant differences due to chance. Further evidence suggesting that these differences are most likely due to chance is the fact that no other liver-function tests besides the total bilirubin showed significant differences and that the greater source of the observed difference in potassium levels was due to a small decline in the placebo group, not a rise in the saw palmetto group (Table 5); the significant difference in glycosuria was due to an increase in urine glucose in placebo-allocated participants. Recent laboratory evidence also suggests that saw palmetto does not have serious hepatic toxicity.²¹

With the growing popularity of dietary supplements, it is imperative that better data on their potential toxicities be generated. Several dietary supplements have been shown to have serious toxic effects and have been removed from the market in the U.S. and some European countries. There is a compelling need to better understand potential adverse effects of other widely used dietary supplements, so that consumers can make more informed decisions about risks and benefits.

The efficacy of saw palmetto extracts for the treatment of BPH is still a matter of controversy and higher-quality studies of this phytotherapeutic are now beginning to appear. Regardless of the ultimate outcomes of these studies, saw palmetto extracts will likely continue to be used widely by men who feel that they benefit from its use.²² Prior studies suggested that side effects of saw palmetto may include headache, dizziness, nausea, and constipation but assessment of adverse effects of saw palmetto has

Table 5 Laboratory test data

Test	Saw palmetto		Placebo		Difference	95% CI	p-Value ^a
	Baseline	12 months	Baseline	12 months			
Albumin (g/dl)	4.17 ± 0.02	4.08 ± 0.03	4.11 ± 0.02	4.04 ± 0.03	-0.01	-0.07 to 0.05	0.44
Bilirubin, total (mg/dl)	0.93 ± 0.03	0.88 ± 0.03	0.90 ± 0.03	0.96 ± 0.03	-0.12	-0.18 to -0.05	0.001
Calcium (mg/dl)	9.64 ± 0.03	9.60 ± 0.03	9.60 ± 0.03	9.59 ± 0.03	-0.04	-0.12 to 0.05	0.44
Carbon dioxide (mmol/dl)	29.59 ± 0.22	28.65 ± 0.22	29.17 ± 0.22	28.73 ± 0.22	-0.50	-1.10 to 0.10	0.20
Chloride (mmol/dl)	101.85 ± 0.27	102.94 ± 0.28	102.22 ± 0.27	103.03 ± 0.28	0.28	-0.52 to 1.09	0.69
Cholesterol, total (mg/dl)	206.22 ± 3.47	205.97 ± 3.53	204.70 ± 3.45	204.02 ± 3.48	0.44	-7.22 to 8.10	0.91
Creatine kinase (CPK) (U/l) ^b	157.06 ± 9.82	154.01 ± 9.97	156.98 ± 9.80	143.19 ± 9.90	10.74	-13.49 to 34.97	0.59
Creatinine (mg/dl)	1.03 ± 0.02	1.03 ± 0.02	1.03 ± 0.02	1.04 ± 0.02	0.001	-0.03 to 0.03	0.46
Glucose (mg/dl)	96.04 ± 2.89	96.08 ± 2.93	97.87 ± 2.87	102.12 ± 2.89	-4.21	-11.19 to 2.77	0.55
Hematocrit (%)	44.34 ± 0.29	43.99 ± 0.29	43.77 ± 0.29	43.91 ± 0.29	-0.48	-1.07 to 0.11	0.43
International normalized ratio (INR) (no units)	0.93 ± 0.01	0.94 ± 0.01	0.92 ± 0.01	0.93 ± 0.01	-0.01	-0.02 to 0.003	0.40
Platelet count (10 ⁹ /l)	241.88 ± 4.85	238.14 ± 4.89	246.83 ± 4.82	237.56 ± 4.85	5.53	-3.16 to 14.22	0.53
Potassium (mmol/dl)	4.32 ± 0.03	4.35 ± 0.04	4.38 ± 0.03	4.33 ± 0.04	0.08	-0.02 to 0.18	0.03
Prostate-specific antigen (PSA) (ng/ml)	1.78 ± 0.14	1.85 ± 0.14	1.62 ± 0.14	1.77 ± 0.14	-0.07	-0.30 to 0.15	0.10
SGOT (AST) (U/l)	26.34 ± 1.23	25.22 ± 1.25	27.80 ± 1.23	28.69 ± 1.24	-2.01	-5.15 to 1.13	0.64
SGPT (ALT) (U/l)	27.28 ± 1.75	25.91 ± 1.77	30.25 ± 1.74	30.82 ± 1.75	-1.94	-5.65 to 1.76	0.56
Sodium (mmol/dl)	138.42 ± 0.21	138.15 ± 0.22	138.43 ± 0.22	138.09 ± 0.22	0.06	-0.62 to 0.75	0.88
Testosterone (ng/dl)	372.97 ± 12.01	356.15 ± 12.10	376.48 ± 11.98	375.07 ± 11.99	-15.40	-39.49 to 8.69	0.06
Triglycerides (mg/dl)	152.97 ± 9.21	143.13 ± 9.32	168.90 ± 9.11	143.62 ± 9.17	15.44	-3.69 to 34.58	0.11
Urea nitrogen (BUN) (mg/dl)	13.92 ± 0.42	14.82 ± 0.43	14.78 ± 0.42	14.75 ± 0.43	0.93	-0.22 to 2.08	0.39
White blood cells (WBC) (10 ⁹ /l)	6.30 ± 0.15	6.08 ± 0.15	5.95 ± 0.15	6.00 ± 0.15	-0.27	-0.60 to 0.05	0.25

Mean laboratory values at baseline and the 12-month closeout visit and between-group differences (calculated as the change in the placebo group subtracted from the change in the saw palmetto group). Mean values are derived from the mixed-effects regression models. The p-values for the between-group differences are derived from the hypothesis test on the group-by-time interaction term from the linear mixed-effects model which tests for overall differences between groups at any time point; the confidence interval for the between-group differences are obtained from the difference between the estimated baseline and closeout values and does not consider interim time points. CI: confidence interval.

^a p-Value for between-group differences in change in variable.

^b Excludes one placebo-randomized participant who developed a statin-induced rhabdomyolysis.

often been incomplete and unsystematic.²³ The STEP trial data are reassuring in that no important toxicities of this extract were identified among the group of patients studied.

These reassuring results, however, must be viewed within the context of the study limitations. The statistical power to detect important clinical differences was limited for some variables, given the sample size of the study. The follow-up phase was 1 year, so no conclusions regarding use over a longer time period can be made. Whether the favorable safety profile of the extract used in this study is typical of other extracts cannot be determined, as there is variation in the extraction techniques and final product composition among the marketed products.²⁴ Finally, rare but serious adverse effects of saw palmetto cannot be assessed in a trial of this size and, like pharmaceutical agents, will require large-scale post-marketing studies to adequately assess this possibility. While case reports do not establish causality, there are case reports suggesting that serious idiosyncratic toxicity of saw palmetto may exist: one patient developed cholestasis after taking an herbal blend that contained saw palmetto,²⁵ another developed transient hepatitis and pancreatitis,²⁶ and one patient suffered excessive intra-operative bleeding and prolonged bleeding time from saw palmetto.²⁷

Overall, the data from the STEP trial do not support the concern of serious clinical adverse effects of this saw palmetto extract over a period of 1 year. While these results are reassuring, further data are needed to more definitively address toxicity issues and will likely emerge from ongoing investigations of saw palmetto as well as population-based toxicity studies.

Conflict of interest statement

None.

Acknowledgments

The authors wish to thank Bertina Lee and Arleen Sakamoto, R.N. for their outstanding assistance during the study, Dr. Henry Leung for management of the study medication, and Dr. Howard Leong for assistance with the laboratory assays. The authors are particularly indebted to the participants without whom this study could not have been conducted.

References

- Committee on the Framework for Evaluating the Safety of the Dietary Supplements National Research Council & Institute of Medicine. Dietary Supplements: A Framework for Evaluating Safety. Executive Summary. 2005 [cited July 9, 2006]; Available from: <http://newton.nap.edu/execsumm.pdf/10882>.
- Lowe FC, Fagelman E. Phytotherapy in the treatment of benign prostatic hyperplasia. *Curr Opin Urol* 2002;12(1):15–8.
- Wilt TJ, Ishani A, Rutks I, MacDonald R. Phytotherapy for benign prostatic hyperplasia. *Public Health Nutr* 2000;3(4A):459–72.
- Guess HA. Epidemiology and natural history of benign prostatic hyperplasia. *Urol Clin North Am* 1995;22(2):247–61.
- Barnes PM, Powell-Griner E, McFann K, Nahin RL. Complementary and alternative medicine use among adults: United States, 2002. *Adv Data* 2004;343:1–19.
- Wilt T, Ishani A, Stark G, Mac Donald R, Mulrow C, Lau J. *Serenoa repens* for benign prostatic hyperplasia (cochrane review). In: *The cochrane library*. Oxford: Update Software; 2002.
- Bent S, Kane C, Shinohara K, Neuhaus J, Hudes ES, Goldberg H, et al. Saw palmetto for benign prostatic hyperplasia. *N Engl J Med* 2006;354(6):557–66.
- Marks LS, Partin AW, Epstein JI, Tyler VE, Simon I, Macairan ML, et al. Effects of a saw palmetto herbal blend in men with symptomatic benign prostatic hyperplasia. *J Urol* 2000;163(5):1451–6.
- Avins AL, Bent S. Saw palmetto and lower urinary tract symptoms: what is the latest evidence? *Curr Urol Rep* 2006;7:260–5.
- Adler SR, Fosket JR. Disclosing complementary and alternative medicine use in the medical encounter: a qualitative study in women with breast cancer. *J Fam Pract* 1999;48(6):453–8.
- Busse JW, Heaton G, Wu P, Wilson KR, Mills EJ. Disclosure of natural product use to primary care physicians: a cross-sectional survey of naturopathic clinic attendees. *Mayo Clin Proc* 2005;80(5):616–23.
- Elder NC, Gillcrist A, Minz R. Use of alternative health care by family practice patients. *Arch Fam Med* 1997;6(2):181–4.
- Eisenberg DM, Kessler RC, Van Rompay MI, Kaptchuk TJ, Wilkey SA, Appel S, et al. Perceptions about complementary therapies relative to conventional therapies among adults who use both: results from a national survey. *Ann Intern Med* 2001;135(5):344–51.
- AARP, National Center for Complementary and Alternative Medicine. Complementary and alternative medicine: What people 50 and older are using and discussing with their physicians. 2007 [cited March 4, 2007]; Available from: <http://assets.aarp.org/rgcenter/health/cam.2007.pdf>.
- Bent S, Padula A, Avins AL. Brief communication: better ways to question patients about adverse medical events. A randomized, controlled trial. *Ann Intern Med* 2006;144(4):257–61.
- O'Leary MP, Fowler FJ, Lenderking WR, Barber B, Sagnier PP, Guess HA, et al. A brief male sexual function inventory for urology. *Urology* 1995;46(5):697–706.
- McCulloch CE, Searle SR. *Generalized, linear and mixed models*. New York: Wiley; 2001.
- Carraro JC, Raynaud JP, Koch G, Chisholm GD, Di Silverio F, Teillac P, et al. Comparison of phytotherapy (Permixon) with finasteride in the treatment of benign prostate hyperplasia: a randomized international study of 1,098 patients. *Prostate* 1996;29(4):231–40. Discussion 41–2.
- Dreikorn K. Phytotherapeutic agents in the treatment of benign prostatic hyperplasia. *Curr Urol Rep* 2000;1(2):103–9.
- Habib FK, Ross M, Ho CK, Lyons V, Chapman K. *Serenoa repens* (Permixon) inhibits the 5alpha-reductase activity of human prostate cancer cell lines without interfering with PSA expression. *Int J Cancer* 2005;114(2):190–4.
- Singh YN, Devkota AK, Sneed DC, Singh KK, Halaweish F. Hepatotoxicity potential of saw palmetto (*Serenoa repens*) in rats. *Phytomedicine* 2007;14(2–3):204–8.
- Blendon RJ, DesRoches CM, Benson JM, Brodie M, Altman DE. Americans' views on the use and regulation of dietary supplements. *Arch Intern Med* 2001;161(6):805–10.
- Wilt TJ, Ishani A, Stark G, MacDonald R, Lau J, Mulrow C. Saw palmetto extracts for treatment of benign prostatic hyperplasia. *JAMA* 1998;280:1604–9.
- Garrard J, Harms S, Eberly LE, Matiak A. Variations in product choices of frequently purchased herbs: caveat emptor. *Arch Intern Med* 2003;163(19):2290–5.

25. Hamid S, Rojter S, Vierling J. Protracted cholestatic hepatitis after the use of prostata. *Ann Intern Med* 1997;127(2):169–70.
26. Jibrin I, Erinle A, Saidi A, Aliyu ZY. Saw palmetto-induced pancreatitis. *South Med J* 2006;99(6):611–2.
27. Cheema P, El-Mefty O, Jazieh AR. Intraoperative haemorrhage associated with the use of extract of saw palmetto herb: a case report and review of literature. *J Intern Med* 2001;250(2):167–9.